

## Recombinant MTF2 Monoclonal Antibody

catalog number: AN302036L

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


|                     |                                                                 |
|---------------------|-----------------------------------------------------------------|
| <b>Reactivity</b>   | Human;Mouse                                                     |
| <b>Immunogen</b>    | Peptide. This information is proprietary to PTMab.              |
| <b>Host</b>         | Rabbit                                                          |
| <b>Isotype</b>      | Rabbit IgG                                                      |
| <b>Clone</b>        | C231657                                                         |
| <b>Purification</b> | Protein A purified                                              |
| <b>Buffer</b>       | PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant. |

### Applications

### Recommended Dilution

|           |        |
|-----------|--------|
| <b>WB</b> | 1:1000 |
|-----------|--------|

### Data



Western Blot with MTF2 Monoclonal Antibody at dilution of 1:1000. Lane 1: HeLa, Lane 2: NIH-3T3

**Observed-MW:47-67 kDa**

**Calculated-MW:67 kDa**

### Preparation & Storage

|                 |                                                                 |
|-----------------|-----------------------------------------------------------------|
| <b>Storage</b>  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| <b>Shipping</b> | Ice bag                                                         |

### Background

MTF2, also named as Metal-response element-binding transcription factor 2, is a 593 amino acid protein. It exists several isoforms and the range of the molecular weight of isoforms are about 55-60 kDa and 29 kDa. MTF2 contains 2 PHD-type zinc fingers and belongs to the Polycomblike family. MTF2 acts by binding to H3K36me3, a mark for transcriptional activation, and recruiting the PRC2 complex, leading to enhance PRC2 H3K27me3 methylation activity and regulates the transcriptional networks during embryonic stem cell self-renewal and differentiation.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.1