

## PE Anti-Mouse CD122 Antibody[TM-Beta 1]

**Catalog Number:** GFH00418D

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

|                                |                                                                                                                                                                                                   |
|--------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Reactivity</b>              | Mouse                                                                                                                                                                                             |
| <b>Host</b>                    | Rat                                                                                                                                                                                               |
| <b>Isotype</b>                 | Rat IgG2b, $\kappa$                                                                                                                                                                               |
| <b>Clone No.</b>               | TM-Beta 1                                                                                                                                                                                         |
| <b>Isotype Control</b>         | PE Rat IgG2b, $\kappa$ Isotype Control[LTF-2] [Product E-AB-F09842D]                                                                                                                              |
| <b>Conjugation</b>             | PE                                                                                                                                                                                                |
| <b>Conjugation Information</b> | PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter). |
| <b>Storage Buffer</b>          | Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide.                                                                                                                               |


### Applications

### Recommended usage

#### FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### Data



Staining of C57BL/6 murine splenocytes cells with APC Anti-Mouse NK1.1 Antibody and PE Anti-Mouse CD122 Antibody[TM-Beta 1] (left) or PE Rat IgG2b,  $\kappa$  Isotype Control (right). Total viable cells were used for analysis.

### Preparation & Storage

|                 |                                                                                                                                                           |
|-----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Storage</b>  | Keep as concentrated solution.<br>This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze. |
| <b>Shipping</b> | Ice bag                                                                                                                                                   |

### Antigen Information

|                        |                                           |
|------------------------|-------------------------------------------|
| <b>Alternate Names</b> | IL-2 Receptor $\beta$ chain;IL-2R $\beta$ |
| <b>Uniprot ID</b>      | P16297                                    |
| <b>Gene ID</b>         | 16185                                     |

### For Research Use Only

## Background

CD122 is a 70-75 kD IL-2 receptor  $\beta$  chain also known as IL-2R $\beta$ , which is also shared by the IL-15 receptor. It is constitutively expressed by NK cells and at lower levels by T cells, B cells, monocytes, and macrophages. The IL-2R $\beta$  chain can combine with either the common  $\gamma$  subunit ( $\gamma_c$ , CD132) alone or with the  $\gamma_c$  subunit and the IL-2R $\alpha$  subunit (CD25) to generate intermediate or high affinity IL-2 receptor complexes, respectively. CD122 expression levels can be upregulated by activation. The TM- $\beta$ 1 antibody does inhibit IL-2 binding to the IL-2 receptor. CD122 is expressed on murine, but not human, CD8+ Tregs involved in the maintenance of T cell homeostasis.