

## Cleaved-PARP1 (D214) Polyclonal Antibody

catalog number: **E-AB-30080**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

<b>Reactivity</b>	Human;Mouse
<b>Immunogen</b>	Synthesized peptide derived from the Internal region of human PARP-1
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Affinity purification
<b>Buffer</b>	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein protectant and 50% glycerol.

### Applications

### Recommended Dilution


<b>WB</b>	1:500-2000
<b>IHC</b>	1:50-300
<b>IF</b>	1:50-300

### Data




Western Blot analysis of A549 cells using Cleaved-PARP1 (D214) Polyclonal Antibody at dilution of 1:2000.

**Observed-MV:24 kDa**  
**Calculated-MV:113 kDa**



Immunofluorescence analysis of Rat spleen tissue using Cleaved-PARP1 (D214) Polyclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using Cleaved-PARP1 (D214) Polyclonal Antibody at dilution of 1:200.

### Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. A void freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

### Background

### For Research Use Only

This gene encodes a chromatin-associated enzyme, poly(ADP-ribosyl)transferase, which modifies various nuclear proteins by poly(ADP-ribosyl)ation. The modification is dependent on DNA and is involved in the regulation of various important cellular processes such as differentiation, proliferation, and tumor transformation and also in the regulation of the molecular events involved in the recovery of cell from DNA damage. In addition, this enzyme may be the site of mutation in Fanconi anemia, and may participate in the pathophysiology of type I diabetes.

## For Research Use Only