

Rat PAI1 Antibody Pair Set

Catalog No.	E-KAB-0644	Applications	ELISA
Synonyms	Endothelial Plasminogen Activator Inhibitor;PAI;PAI-1;PLANH1;SERPINE1		

Kit components & Storage

Title	Specifications	Storage
Rat PAI1 Capture Antibody	1 vial, 100 µg	Store at -20℃ for one year. Avoid freeze/thaw cycles.
Rat PAI1 Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20℃ for one year. Avoid freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Product Information

Items		Characteristic (E-KAB-0644)	
		Rat PAI1 Capture Antibody	Rat PAI1 Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Rat PAI1 protien	Recombinant Rat PAI1 protien
	Swissprot	P20961	
Product details	Reactivity	Rat	Rat
	Host	Rabbit	Rabbit
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50% glycerol; pH 7.5	PBS with 0.04% Proclin 300; 1% protective protein; 50% glycerol; pH 7.5
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Rat PAI1 in ELISAs.	

For Research Use Only

Applications

Rat PAI1 Sandwich ELISA Assay

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4 µg/mL	Rat PAI1 Capture Antibody	
ELISA Detection	1:1000-1:10000	Rat PAI1 Detection Antibody (Biotin)	

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Serine protease inhibitor. Inhibits TMPRSS7. Is a primary inhibitor of tissue-type plasminogen activator (PLAT) and urokinase-type plasminogen activator (PLAU) . As PLAT inhibitor , it is required for fibrinolysis down-regulation and is responsible for the controlled degradation of blood clots. As PLAU inhibitor , it is involved in the regulation of cell adhesion and spreading. Acts as a regulator of cell migration , independently of its role as protease inhibitor. It is required for stimulation of keratinocyte migration during cutaneous injury repair. It is involved in cellular and replicative senescence. Plays a role in alveolar type 2 cells senescence in the lung. Is involved in the regulation of cementogenic differentiation of periodontal ligament stem cells , and regulates odontoblast differentiation and dentin formation during odontogenesis.

For Research Use Only