Recombinant Human Amyloid Precursor (C-FC)

Catalog Number: PKSH033887

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	HEK293 Cells-derived Human Amyloid Precursor protein Leu18-Lys612, with an C-
	terminal Fc
Calculated MW	94.6 kDa
Observed MW	120-145 kDa
Accession	P05067-4
Bio-activity	Not validated for activity
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}C$ for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, 5% Trehalose, pH7.4.
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Web:www.elabscience.com

Elabscience®

Amyloid precursor protein (APP) is a type I membrane protein with several isoforms due to alternative splicing, performs physiological functions on the surface of neurons relevant to neurite growth, neuronal adhesion and axonogenesis. Of the three major splice isoforms of APP (APP695, APP751, and APP770) APP695 is the predominant neuronal form from which Amyloid beta peptide and transcriptionally-active cleaved intracellular domain of APP (AICD) are preferentially generated by selective processing through the amyloidogenic pathway. Human APP695 consists of a 17 amino acid (aa) signal sequence, a 607 aa extracellular domain (ECD), a 24 aa transmembrane domain, and a 47 aa cytoplasmic domain. Within the ECD, human APP695 shares 97% aa sequence identity with mouse and rat APP695. Amyloid beta is a major molecule implicated in pathogenesis of Alzheimer's disease (AD) and related dementias. AICD regulates expression by direct promoter binding of multiple genes, including APP itself, the beta-secretase, BACE-1 and the Amyloid beta-degrading enzyme, Neprilysin. As such, APP695 plays an important role in brain development, learning and memory, synaptic plasticity, and neurodegeneration including AD.