

Recombinant Mouse CD16/FCGR3 Protein (His Tag)

Catalog Number: PKSM041338

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse CD16/FCGR3 protein Leu32-Thr215, with an C-terminal His
Calculated MW	21.9 kDa
Observed MW	35-40 kDa
Accession	Q5D5I8
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Low affinity immunoglobulin gamma Fc region receptor III (Fc gamma RIII/CD16) is a member of the Ig superfamily. Based on close relationships in their extracellular domains, the Fc gamma Rs have been divided into three classes composing of Fc gamma RI (CD64), Fc gamma RII (CD32), and Fc gamma RIII (CD16). Each group may be encoded by multiple genes and exist in different isoforms depending on species and cell type. Mouse CD16 is a type I transmembrane protein having two extracellular Ig-like domains consisting of immunoglobulin domain, repeat, signa and transmembrane, transmembrane helix. It is expressed on a variety of myeloid and lymphoid cells and associates with Fc R gamma to deliver an activating signal upon ligand binding. Fcgr3 is IgG binding and activation or inhibition of immune responses such as antibody-dependent cellular cytotoxicity, phagocytosis, cell surface receptor signaling pathway and positive regulation of type I/IIa/III hypersensitivity.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017