

# Purified Anti-Mouse CD71 Antibody[R17 217.1.3/TIB-219]

catalog number: E-AB-F1093A



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description


<b>Reactivity</b>	Mouse
<b>Immunogen</b>	Recombinant Mouse CD71 protein
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG2a, $\kappa$
<b>Clone</b>	R17 217.1.3/TIB-219
<b>Purification</b>	>98%, Protein A/G purified
<b>Conjugation</b>	Unconjugated
<b>buffer</b>	PBS, pH 7.2. Contains 0.05% proclin 300.

## Applications

## Recommended Dilution

<b>FCM</b>	2 $\mu\text{g/mL}$ ( $1 \times 10^5$ - $5 \times 10^5$ cells)
------------	---

## Data



C57/BL6 Mouse bone marrow cells were stained with 0.2  $\mu\text{g}$  AF/LE Purified Anti-Mouse CD71 Antibody[R17 217.1.3/TIB-219](Right) and 0.2  $\mu\text{g}$  Rat IgG2a,  $\kappa$  Isotype Control(Left), followed by Alexa Fluor<sup>®</sup> 647-conjugated Goat Anti-Rat IgG Secondary Antibody, then anti-Mouse Ter-119 PE-conjugated Monoclonal Antibody.

## Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	Ice bag

## Background

CD71 is a 95 kD type II heterodimeric transmembrane glycoprotein that is also known as T9 and transferrin receptor. CD71 is expressed on proliferating cells, reticulocytes, and erythroid precursors. Its expression is very low on resting leukocytes. CD71 plays a role in the control of cellular proliferation by facilitating the uptake of iron via ferrotransferrin binding and the recycling of apotransferrin to the cell surface.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine  
Tel: 400-999-2100

Email: [techsupport@elabscience.cn](mailto:techsupport@elabscience.cn)

Web: [www.elabscience.cn](http://www.elabscience.cn)

Rev. V2.3