



A Reliable Research Partner in Life Science and Medicine

# **Mouse IL-10 Antibody Pair Set**

Catalog No. E-KAB-0079 Applications ELISA

**Synonyms** IL10, IL10A, CSIF, TGIF

## **Kit components & Storage**

Title	Specifications	Storage
Mouse IL-10 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Mouse IL-10 Detection Antibody (Biotin)	1 vial, 50 μL	Store at -20℃ for one year.
		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0079)	
		Mouse IL-10 Capture Antibody	Mouse IL-10 Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Mouse IL-10 protein	Recombinant Mouse IL-10 protein
Information	Swissprot	P18893	
Product details	Reactivity	Mouse	Mouse
	Host	Rat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Mouse IL-10 in ELISAs.	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>



A Reliable Research Partner in Life Science and Medicine

### **Applications**

Mouse IL-10 Sandwich ELISA Assav:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Mouse IL-10 Capture Antibody	
Capture			Africa Af
ELISA Detection	1:1000-1:10000	Mouse IL-10 Detection Antibody (Biotin)	0.01 100 1000 10000 Mouse IL-10 concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

# **Background**

This gene encodes an anti-inflammatory cytokine that is a member of the class-2 cytokine family. The encoded protein is secreted by cells of both the innate and adaptive immune systems and is crucial for limiting the immune response to a broad range of pathogens. It also has been shown to suppress autoimmune responses. This protein mediates it's immunosuppressive signal through a specific interleukin 10 receptor complex. Aberrant functioning of this gene is associated with numerous immune disorders including graft-versus-host disease, and increased susceptibility to HIV-1 infection and rheumatoid arthritis.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>