



A Reliable Research Partner in Life Science and Medicine

Recombinant Human LMNA Protein(Gst Tag)

Catalog Number: PDEH100526

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| \mathbf{r} | | | | | | | |
|--------------|----|---|---|---|---|----|---|
| H) | es | C | m | n | т | ſΠ | ï |

Species Human

Source E.coli-derived Human LMNA protein Met1-Ser212, with an N-terminal GST

 Mol_Mass
 49.2 kDa

 Accession
 P02545

Bio-activity Not validated for activity

Properties

Purity > 90% as determined by reducing SDS-PAGE.

Endotoxin < 10 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human LMNA proteins, 2 µg/lane of Recombinant Human LMNA proteins was resolved with an SDS-PAGE under reducing conditions, showing bands at 49.2 KD

Background

The nuclear lamina consists of a two-dimensional matrix of proteins located next to the inner nuclear membrane. The lamin family of proteins make up the matrix and are highly conserved in evolution. During mitosis, the lamina matrix is reversibly disassembled as the lamin proteins are phosphorylated. Lamin proteins are thought to be involved in nuclear stability, chromatin structure and gene expression. Vertebrate lamins consist of two types, A and B. Alternative splicing results in multiple transcript variants. Mutations in this gene lead to several diseases: Emery-Dreifuss muscular dystrophy, familial partial lipodystrophy, limb girdle muscular dystrophy, dilated cardiomyopathy, Charcot-Marie-Tooth disease, and Hutchinson-Gilford progeria syndrome.

For Research Use Only