

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® Violet 540 Anti-Human CD19 Antibody[CB19]

Catalog Number: E-AB-F1004T3

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human;Rhesus;Cynomolgus

Host Mouse

Isotype Mouse IgG1, κ

Clone No. CB19

Isotype Control Elab Fluor® Violet 540 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-

F09792T31

Conjugation Elab Fluor® Violet 540

Conjugation Information Elab Fluor® Violet 540 is designed to be excited by the violet laser (405 nm) and detected

using an optical filter centered near 548 nm (e.g., a 572/28 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



Staining of normal human peripheral blood cells with Elab

Fluor[®] Violet 540 Anti-Human CD19 Antibody[CB19] (filled gray histogram) or Elab Fluor[®] Violet 540 Mouse IgG1, κ Isotype Control(empty black histogram). Cells in the lymphocytes gate were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names B-lymphocyte antigen CD19;CD19;Cd19;Differentiation antigen CD19

Uniprot ID P15391

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Gene ID Background 930

CD19 is a 95 kD type I transmembrane glycoprotein also known as B4. It is a member of the immunoglobulin superfamily expressed on B-cells (from pro-B to blastoid B cell s, absent on plasma cells) and follicular dendritic cells. CD19 is involved in B cell development, activation, and differentiation. CD19 forms a complex with CD21 (CR2) and CD81 (TAPA-1), and functions as a BCR co-receptor.