



A Reliable Research Partner in Life Science and Medicine

Human IL-15 Antibody Pair Set

Catalog No. E-KAB-0704 Applications ELISA

Synonyms IL15;IL-15;IL-15MGC9721;interleukin 15;interleukin-15

Kit components & Storage

Title	Specifications	Storage
Human IL-15 Capture Antibody	1 vial, 100 μ g	Store at -20°C. Avoid freeze / thaw
		cycles.
Human IL-15 Detection Antibody	1 vial, 50 μL	Store at -20°C. Avoid freeze / thaw
(Biotin)		cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0704)		
		Human IL-15 Capture Antibody	Human IL-15 Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Human IL-15 protein	Recombinant Human IL-15 protein	
Information	Swissprot	P40933		
Product details	Reactivity	Human	Human	
	Host	Mouse	Mouse	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
			7.4	
	Purify	Protein A or G	Protein A or G	
	Specificity	Detects Human IL-15 in ELISAs.		

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com



A Reliable Research Partner in Life Science and Medicine

Applications

Human IL-15 Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4ug/mL	Human IL-15 Capture	
Capture		Antibody	10
			Optical Density
ELISA	1:1000-1:10000	Human IL-15 Detection	Optical Optical
Detection		Antibody (Biotin)	•
			0.01 10 100 1000 10000 Human IL-15 Concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

IL-15 is a regulatory cytokine, and it is produced by dendritic cells, epithelial cells, fibroblasts, and monocytes. IL-15 plays an important role in immune response and shares many functions with IL-2. For example, it stimulates the proliferation of activated T cells, NK cells, and B cells, and induces immunoglobulin synthesis by B cells stimulated by anti-IgM or CD40 ligand. In addition, IL-15 promotes the development of dendritic cells and induces the production of proinflammatory cytokines from macrophages. IL-15 acts as a bridge between innate and adaptive immunity because of its diverse roles in the immune system.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com