



## FAM111B Polyclonal Antibody

catalog number: E-AB-19841

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

| Description  |  |
|--------------|--|
| Reactivity   | Human  |
| Immunogen    | Synthetic peptide of human FAM111B   |
| Host         | Rabbit   |
| Isotype      | IgG  |
| Purification | Antigen affinity purification  |
| Conjugation  | Unconjugated   |
| Buffer       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilution   |
| IHC          | 1:50-1:300   |

| Data  |   |
|---|---|
|   |    |
| Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using FAM111B Polyclonal Antibody at dilution of 1:50(×200) | Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using FAM111B Polyclonal Antibody at dilution of 1:50(×200) |

| Preparation & Storage |  |
|-----------------------|--|
| Storage               | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.  |
| Shipping              | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

| Background   |  |
|--|--|
| <p>This gene encodes a protein with a trypsin-like cysteine/serine peptidase domain in the C-terminus. Mutations in this gene are associated with an autosomal dominant form of hereditary fibrosing poikiloderma (HFP). Affected individuals display mottled pigmentation, telangiectasia, epidermal atrophy, tendon contractures, and progressive pulmonary fibrosis. Alternative splicing results in multiple transcript variants encoding distinct isoforms. A paralog of this gene which also has a trypsin-like peptidase domain, FAM111A, is located only 16 kb from this gene on human chromosome 11q12.1.</p> |  |