

IL-5/EDF/BCDFII/TRF, Mouse, Recombinant**Cat. No. : GPCK315****产品信息**

物种	Mouse
表达宿主	E.coli
序列信息	MEIPMSTVVKETLTQLSAHRALLTSNETMRLPVPVTHKNHQLCIGEIFQGLDILKNQTVRGGTVE MLFQNLSLIKKYIDRQKEKCGEERRRTRQFLDYLQEFLGVMSTEWAMEG with polyhistidine tag at the C-terminus.
检索号	P04401.1
标签	His-tag at the C-terminus
分子量	13.93 kDa
有效期	12 months
生物活性	Measure by its ability to induce TF-1 cells proliferation. The ED50 for this effect is < 0.2 ng/mL. The specific activity of recombinant mouse IL-5 is > 5 × 10 ⁶ IU/mg.

产品特性

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 μm filtered solution containing 1 × PBS, pH 7.4.
复溶	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Interleukin 5 (IL-5) is an interleukin produced by type-2 T helper cells and mast cells. Through binding to the interleukin-5 receptor, interleukin 5 stimulates B cell growth and increases immunoglobulin secretion-primarily IgA. It is also a key mediator in eosinophil activation.

