A Reliable Research Partner in Life Science and Medicine

# Recombinant Rat IL-2 Protein(His Tag)

Catalog Number: PDER100250

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

**Species** Rat

Source E.coli-derived Rat IL-2 protein Ala21-Gln155, with an N-teminal His

Calculated MW14.7 kDaObserved MW14 kDaAccessionP17108

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 95% as determined by reducing SDS-PAGE.

**Endotoxin** < 10 EU/mg of the protein as determined by the LAL method

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

**Reconstitution** It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



SDS-PAGE analysis of Rat IL-2 proteins, 2µg/lane of Recombinant Rat IL-2 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 14 kDa

## Background

Interleukin-2, also known as a T-cell growth factor, TCGF, Aldesleukin, and II.2, is a secreted protein that belongs to the IL-2 family. Interleukin-2 / IL-2 was the first interleukin molecule to be discovered. Interleukin-2 / IL-2 molecule was first purified to homogeneity by immunoaffinity chromatography by Kendall Smith and his team at Dartmouth Medical Schoo l. Interleukin-2 / IL-2 was also the first cytokine shown to mediate its effects via a specific IL-2 receptor, and it was also the first interleukin to be cloned and expressed from a complementary DNA (cDNA) library. Interleukin-2 / IL-2 was designated number 2 because Smith's data at the time indicated that IL-1, produced by macrophages, facilitates IL-2 production by T lymphocytes (T cells).

### For Research Use Only