



A Reliable Research Partner in Life Science and Medicine

# **MCM7 Polyclonal Antibody**

catalog number: E-AB-40231

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Reactivity Human

Immunogen Recombinant Human DNA replication licensing factor MCM7 protien

Host Rabbit
Isotype IgG


**Purification** Antigen Affinity Purification

**Buffer** PBS with 0.05% proclin 300, 1% protective protein and 50% glycerol,pH7.4

## **Applications** Recommended Dilution

**IHC** 1:50-1:200

#### Data



Immunohistochemistry of paraffin-embedded Human lymph node using MCM7 Polyclonal Antibody at dilution of 1:200

### Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

# Background

The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance proteins (MCM) that are essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex (pre\_RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1-dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumorsuppressor protein RB1/RB. Alternatively spliced transcript variants encoding distinct isoforms have been reported.