

## Phospho-AMPK alpha1/2 (Thr183/172) Polyclonal Antibody

catalog number: E-AB-21121

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human; Mouse; Rat; Monkey

**Immunogen** Synthesized peptide derived from human AMPK $\alpha$ 1/2 around the phosphorylation site

of Thr183/172

**Host** Rabbit Isotype IgG

**Purification** Affinity purification


**Buffer** Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein

protectant and 50% glycerol.

# Applications Recommended Dilution WB 1:500-1:2000

IHC 1:100-1:300 IF 1:50-1:200


#### Data



Western Blot analysis of various cells using Phospho-AMPK alpha1/2 (Thr183/172) Polyclonal Antibody at dilution of 1:500

Immunohistochemistry of paraffin-embedded mouse brain using Phospho-AMPK alpha1/2 (Thr183/172) Polyclonal Antibody at dilution of 1:50

### Observed-MW:63 kDa Calculated-MW:62 kDa



Immunofluorescence analysis of Rat heart tissue using Phospho-AMPK alpha1/2 (Thr183/172) Polyclonal Antibody at dilution of 1:200

#### Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

#### Background

#### For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com Fax: 1-832-243-6017

# Elabscience®

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

AMPK (for 5'-AMP-activated protein kinase) is a heterotrimeric complex comprising a catalytic  $\alpha$  subunit and regulatory  $\beta$  and  $\gamma$  subunits. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. AMPK is activated by high AMP and low ATP through a mechanism involving allosteric regulation, promotion of phosphorylation by an upstream protein kinase known as AMPK kinase, and inhibition of dephosphorylation. Activated AMPK can phosphorylate and regulate in vivo hydroxymethylglutaryl-CoA reductase and acetyl-CoA carboxylase, which are key regulatory enzymes of sterol synthesis and fatty acid synthesis, respectively

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com