Recombinant Mouse Ceacam1 Protein(His Tag)


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Catalog Number: PDMM100233



Description **Species** Mouse Source Mammalian-derived Mouse Ceacam1 protein Glu35-Gly428, with an C-terminal His Mol Mass 43.2 kDa Accession P31809 **Bio-activity** Not validated for activity **Properties** Purity >90% as determined by reducing SDS-PAGE. Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. This product is provided as lyophilized powder which is shipped with ice packs. Shipping Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Formulation Mannitol. Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis

Data



SDS-PAGE analysis of Mouse Ceacam1 proteins, 2µg/lane of Recombinant Mouse Ceacam1 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 55-80 kDa

Background

For Research Use Only

Recombinant Mouse Ceacam1 Protein(His Tag)



Catalog Number: PDMM100233

The carcinoembryonic-antigen-related cell-adhesion molecule (CEACAM) family of proteins has been implicated in various intercellular-adhesion and intracellular-signalling-mediated effects that govern the growth and differentiation of normal and cancerous cells. CEACAM1, also known as biliary glycoprotein I (BGP I) and CD66a, is a member of the carcinoembryonic antigen (CEA) gene family which belongs to the immunoglobulin superfamily. The highly glycosylated CEACAM1 contains one N-terminal V-type Ig-like domain and three C2-type Ig-like domains within its EC D, and one ITIM motif and a calmodulin binding site in the cytoplasmic region. CEACAM1 is a surface glycoprotein expressed on various blood cells, epithelial cells, and vascular cells. It was described as an adhesion molecule mediating cell adhesion via both homophilic and heterophilic manners, and was detected on leukocytes, epithelia, and endothelia. Studies have revealed that CEACAM1 performs actions in multiple cellular processes including tissue differentiation, angiogenesis, apoptosis, metastasis, as well as the modulation of innate and adaptive immune responses.

For Research Use Only