Recombinant Rat IGFBP6/IBP6 Protein (His Tag)

Catalog Number: PKSR030133

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Species	Rat
Source	HEK293 Cells-derived Rat IGFBP6/IBP6 protein Met1-Gly226, with an C-terminal His
Calculated MW	22.9 kDa
Accession	NP_037236.1
Bio-activity	Measured by its ability to inhibit the biological activity of IGFII on MCF7 human breast adenocarcinoma cells. The ED_{50} for this effect is typically 0.6-3 μg/mL in the presence of 14 ng/mL human IGFII.

Properties

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



Measured by its ability to inhibit the biological activity of IGFII on MCF7 human breast adenocarcinoma cells. The ED_{50} for this effect is typically 0.6-3 μg/mL in the presence of 14 ng/mL human IGFII.

Background

Insulin-like growth factor binding protein 6 (IGFBP6) is a 24-kDa protein that binds insulin-like growth factor 1 (IGF-1) and IGF-2 with high affinity and inhibits IGF action in vitro. The Insulin-like growth factor-binding protein also known as IGFBP serves as a carrier protein for Insulin-like growth factor 1. IGFBPs are clearly distinct but are sharing regions with strong homology. All members of the IGFBP family bind IGF-I and IGF-II with about equal affinity. Insulin-like growth factor (IGF) binding proteins (IGFBPs) have been shown to either inhibit or enhance the action of IGF, or act in an IGF-independent manner in the prostate. IGF-binding protein-4 (IGFBP-4) inhibits IGF-I action in vitro and is the most abundant IGFBP in the rodent arterial wall. IGFBP6 is directly downregulated by the beta-catenin/TCF complex in desmoid tumors, and imply a role for the IGF axis in the proliferation of desmoid tumors. There is mounting evidence that the structure of the IGFBP proteins plays a key role in the regulation of IGF bioavailability, by modulating its molecular size, capillary membrane permeability, target tissue specificity, cell membrane adherence and IGF affinity.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com