
## Histone H2A.X Polyclonal Antibody


#### catalog number: E-AB-70233

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	KLH conjugated Synthetic peptide corresponding to Mouse Histone H2A.X
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 1% protein
	protectant and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:300-1:800


#### Data





Western Blot analysis of various samples using Histone H2A.X Polyclonal Antibody at dilution of 1:1000.

### Observed-MW:18 kDa Calculated-MW:15-18 kDa



testis using Histone H2A.X Polyclonal Antibody at dilution of 1:400.

Immunohistochemistry analysis of paraffin-embedded rat testis using Histone H2A.X Polyclonal Antibody at dilution of 1:400.

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

#### For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com Fax: 1-832-243-6017

# **Elabscience**®

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif.

For Research Use Only