# IL-18/IL-1F4 Polyclonal Antibody(Capture/Detector)

catalog number: AN000230P



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| -            |   |   |    |    |    | ٠, |    |   |
|--------------|---|---|----|----|----|----|----|---|
| D            | Α |   | C  | PТ | m  | ŤΤ | U. | m |
| $\mathbf{L}$ | v | 0 | v. | ш  | עו | UΙ | U. |   |

**Reactivity** Human

Immunogen Recombinant Human IL-18/IL-1F4 protein expressed by E.coli

Host Rabbit
Isotype Rabbit IgG

**Purification** Antigen Affinity Purification


**Conjugation** Unconjugated

buffer Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

Applications Recommended Dilution

**ELISA Capture** 2-8 μg/mL **ELISA Detector** 0.1-0.4 μg/mL

#### Data



Sandwich ELISA-Recombinant Human IL-18/IL-1F4 protein standard curve. Background subtracted standard curve using IL-18/IL-1F4 antibody(AN000230P)(Capture), IL-18/IL-1F4 antibody(AN000230P)(Detector) in sandwich ELISA. The reference range value for Recombinant Human IL-18/IL-1F4 protein is 31.25-2000 pg/mL.

#### **Preparation & Storage**

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

### Background

Pro-inflammatory cytokine primarily involved in epithelial barrier repair, polarized T-helper 1 (Th1) cell and natural killer (NK) cell immune responses. Upon binding to II18R1 and II18RAP, forms a signaling ternary complex which activates NF-kappa-B, triggering synthesis of inflammatory mediators. Synergizes with II12/interleukin-12 to induce IFNG synthesis from T-helper 1 (Th1) cells and natural killer (NK) cells. Involved in transduction of inflammation downstream of pyroptosis: its mature form is specifically released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore.

## For Research Use Only