



A Reliable Research Partner in Life Science and Medicine

Human ApoA1 Antibody Pair Set

Catalog No. E-KAB-0260 Applications ELISA

Synonyms APOA1, Apo-A1

Kit components & Storage

Title	Specifications	Storage
Human ApoA1 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Human ApoA1 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0260)		
		Human ApoA1 Capture Antibody	Human ApoA1 Detection Antibody (Biotin)	
Immunogen	Immunogen	Recombinant Human ApoA1 protein	Recombinant Human ApoA1 protein	
Information	Swissprot	P02647		
Product details	Reactivity	Human	Human	
	Host	Mouse	Mouse	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
			7.4	
	Purify	Protein A	Protein A	
	Specificity	Detects Human ApoA1 in ELISAs.		

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com



Elabscience®

A Reliable Research Partner in Life Science and Medicine

Applications

Human ApoA1 Sandwich ELISA Assav:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Human ApoA1 Capture Antibody	
Capture			10] Ag
ELISA Detection	1:1000-1:10000	Human ApoA1 Detection Antibody (Biotin)	Optical Density
			0.01 1 10 100 1000 Human ApoA1 concentration(ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

ApoA1 is a major protein component of high density lipoproteins (HDL) which is associated with reversed cholesterol transport, lipid/cholesterol binding, lecithin/cholesterol acyltransferase (LCAT) activation and specific receptors binding. It is synthesized in the liver and small intestine. Defects of ApoA1 cause low HDL level and systemic non-neuropathic amyloidosis. Serum concentration of ApoA1 is inversely related to the risk of developing atherosclerosis. This antibody was generated against the C-terminal region of human ApoA1.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com