A Reliable Research Partner in Life Science and Medicine

# Recombinant Human Protamine-2/PRM2 Protein (His Tag)

Catalog Number: PKSH032952

Note: Centrifuge before opening to ensure complete recovery of vial contents.

$\mathbf{r}$							
H)	es	C	m	n	т	ſΠ	ï

Species Human

Source HEK293 Cells-derived Human Protamine-2;PRM2 protein Met 1-Phe389, with an C-

terminal His

Calculated MW45.9 kDaObserved MW47 kDaAccessionP31350

**Bio-activity** Not validated for activity

#### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, 5%

Trehalose, pH 8.0.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Ribonucleoside-Diphosphate Reductase Subunit M2 (RRM2) belongs to the ribonucleoside diphosphate reductase small chain family. The reductase of RRM2 catalyzes the formation of deoxyribonucleotides from ribonucleotides. Synthesis of the encoded protein (M2) is regulated in a cell-cycle dependent fashion. RRM2 supplies the precursors essential for DNA synthesis. RRM2 catalyzes the biosynthesis of deoxyribonucleotides from the corresponding ribonucleotides. Phosphorylation on Ser-20 relieves the inhibitory effect on Wnt signaling.

#### For Research Use Only