

## AF/LE Purified Anti-Mouse IL-17A Antibody[17F3]

**Catalog Number:** GF12720

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


|                     |   |
|---------------------|---|
| <b>Reactivity</b>   | Mouse   |
| <b>Immunogen</b>    | Recombinant Mouse IL-17A protein  |
| <b>Host</b>         | Mouse   |
| <b>Isotype</b>      | Mouse IgG1, κ   |
| <b>Clone</b>        | 17F3  |
| <b>Purification</b> | >98%, Protein A/G purified  |
| <b>Conjugation</b>  | None (AF/LE)  |
| <b>Buffer</b>       | Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method. |

### Applications

### Recommended Dilution

|            |  |
|------------|--|
| <b>FCM</b> | 2 µg/mL (0.5×10 <sup>6</sup> -1×10 <sup>6</sup> cells) |
|------------|--|

### Data



HEK293T cells transfected with pcDNA3.1 plasmid encoding Mouse IL-17A gene were stained with 0.2 µg AF/LE Purified Anti-Mouse IL-17A Antibody[17F3] (Right) and 0.2 µg Mouse IgG1, κ Isotype Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody.

### Preparation & Storage

|                 |  |
|-----------------|--|
| <b>Storage</b>  | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions. |
| <b>Shipping</b> | Ice bag  |

### Background

The 17F3 monoclonal antibody reacts with mouse IL-17A a 15-20 kDa cytokine expressed by Th17 cells, γδ T cells, iNKT cells, NK cells, LTI cells, neutrophils, and intestinal Paneth cells. IL-17A has pleiotropic effects in immunoregulation and inflammation. It plays an important role in anti-microbial and chronic inflammation by inducing cytokine and chemokine production, neutrophil influx, and the production of antibacterial peptides but it is also an inflammatory mediator in the development of autoimmune diseases including rheumatoid arthritis, asthma, multiple sclerosis, and psoriasis. The 17F3 antibody has been shown to neutralize IL-18A in vivo.