# Recombinant Human ICAM-1/CD54 Protein

**Catalog Number: PKSH031660**

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*


## Description

| | |
|---|---|
| **Species** | Human |
| **Source** | HEK293 Cells-derived Human ICAM-1/CD54 protein Met 1-Glu 480 |
| **Calculated MW** | 51.0 kDa |
| **Accession** | NP_000192.2 |
| **Bio-activity** | Not validated for activity |

## Properties

| | |
|---|---|
| **Purity** | > 95 % as determined by reducing SDS-PAGE. |
| **Endotoxin** | < 1.0 EU per μg of the protein as determined by the LAL method. |
| **Storage** | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| **Shipping** | This product is provided as lyophilized powder which is shipped with ice packs. |
| **Formulation** | Lyophilized from sterile PBS, pH 7.4. |
| **Reconstitution** | Please refer to the printed manual for detailed information. |

## Data




> 95 % as determined by reducing SDS-PAGE.

## Background

Intercellular adhesion molecule-1 (ICAM-1, or CD54) is a 90 kDa member of the immunoglobulin (Ig) superfamily and is critical for the firm arrest and transmigration of leukocytes out of blood vessels and into tissues. ICAM-1 is constitutively present on endothelial cells, but its expression is increased by proinflammatory cytokines. The endothelial expression of ICAM-1 is increased in atherosclerotic and transplant-associated atherosclerotic tissue and in animal models of atherosclerosis. Additionally, ICAM-1 has been implicated in the progression of autoimmune diseases. ICAM-1 is a ligand for LFA-1(integrin). When activated, leukocytes bind to endothelial cells via ICAM-1/LFA-1 interaction and then transmigrate into tissues. Presence with heavy glycosylation and other structural characteristics, ICAM-1 possesses binding sites for a number of immune-associated ligands and serves as the binding site for entry of the major group of human Rhinovirus (HRV) into various cell types. ICAM-1 also becomes known for its affinity for Plasmodium falciparum-infected erythrocytes (PFIE), providing more of a role in infectious disease. Previous studies have shown that ICAM-1 is involved in inflammatory reactions and that a defect in ICAM-1 gene inhibits allergic contact hypersensitivity.


Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com