

Recombinant Mouse CCL11 Protein(Trx Tag)

Catalog Number: GPEM0147

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	E.coli-derived Mouse CCL11 protein His24-Pro97, with an N-terminal Trx
Calculated MW	28.1 kDa
Observed MW	32 kDa
Accession	P48298
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Mouse CCL11 proteins, 2 µg/lane of Recombinant Mouse CCL11 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 28.1 KD

Background

CCL11 or chemokine (C-C motif) ligand 11 is a member of the chemokine (C-C motif) ligand family. Chemokine (C-C motif) ligand 11 is a member of the chemokine family. There are four members of the chemokine family: C-C chemokines, C chemokines, CXC chemokines and CX3C chemokines. The C-C chemokines have two cysteines nearby the amino terminus. There have been at least 27 distinct members of this subgroup reported for mammals, called C-C chemokine ligands (CCL)-1 to 28. Chemokines are a family of small chemotactic cytokines, or proteins secreted by cells. They share the same structure similarities such as small size, and the presence of four cysteine residues in conserved locations in order to form their 3-dimensional shape. Some of the chemokines are considered pro-inflammatory which can be induced to recruit cells of the immune system to a site of infection during an immune response, while others are considered homeostatic and are implied in controlling the migration of cells during normal processes of tissue maintenance and development. CCL11 is implicated in allergic responses through selectively recruiting eosinophils by inducing their chemotaxis. The effects of CCL11 are mediated by its binding to chemokine receptor. Increased CCL11 levels in blood plasma are associated with aging in mice.