

## Recombinant VEGF Receptor 1 Monoclonal Antibody

catalog number: **AN301010L**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

|                     |   |
|---------------------|---|
| <b>Reactivity</b>   | Human;Mouse;Rat   |
| <b>Immunogen</b>    | Recombinant Human VEGF Receptor 1 protein                       |
| <b>Host</b>         | Rabbit  |
| <b>Isotype</b>      | IgG, $\kappa$   |
| <b>Clone</b>        | B761  |
| <b>Purification</b> | Protein A   |
| <b>Buffer</b>       | PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant. |

### Applications


| Applications | Recommended Dilution |
|--------------|----------------------|
| IHC          | 1:200-1:1000         |
| WB           | 1:1000-1:5000        |

### Data



Western Blot with Recombinant VEGF Receptor 1 Monoclonal Antibody at dilution of 1:1000 dilution. Lane A: K562 cells.

**Observed-MW:180 kDa**  
**Calculated-MW:151 kDa**



Immunohistochemistry of paraffin-embedded human stomach carcinoma tissue using Recombinant VEGF Receptor 1 Monoclonal Antibody at dilution of 1:200.

### Preparation & Storage

|                 |   |
|-----------------|---|
| <b>Storage</b>  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| <b>Shipping</b> | Ice bag   |

### Background

FLT1(fms related tyrosine kinase 1) Homo sapiens This gene encodes a member of the vascular endothelial growth factor receptor (VEGFR) family. VEGFR family members are receptor tyrosine kinases (RTKs) which contain an extracellular ligand-binding region with seven immunoglobulin (Ig)-like domains, a transmembrane segment, and a tyrosine kinase (TK) domain within the cytoplasmic domain. This protein binds to VEGFR-A, VEGFR-B and placental growth factor and plays an important role in angiogenesis and vasculogenesis. Expression of this receptor is found in vascular endothelial cells, placental trophoblast cells and peripheral blood monocytes. Multiple transcript variants encoding different isoforms have been found for this gene. Isoforms include a full-length transmembrane receptor isoform and shortened, soluble isoforms. The soluble isoforms are associated with the onset of pre-eclampsia.

### For Research Use Only