

Recombinant Human IL18 C terminal protein (Avi,His Tag)

Catalog Number: PDEH100868



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	E.coli-derived Human IL18 C terminal protein Ile116-Asp193, with an C-terminal Avi & His
Mol_Mass	8.5 kDa
Accession	Q14116
Bio-activity	Not validated for activity

Properties

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

A Reliable Research Partner in Life Science and Medicine
Tel:400-999-2100

Email:techsupport@elabscience.cn

Web:www.elabscience.cn

Rev. V1.3

Recombinant Human IL18 C terminal protein (Avi,His Tag)

Catalog Number: PDEH100868



Interleukin-18 (IL-18, also known as interferon-gamma inducing factor) is a proinflammatory cytokine that belongs to the IL-1 superfamily and is produced by macrophages and other cells. This cytokine can induce the IFN-gamma production of T cells. The combination of IL-18 and IL12 has been shown to inhibit IL4 dependent IgE and IgG1 production, and enhance IgG2a production of B cells. IL-18 binding protein (IL18BP) can specifically interact with this cytokine, and thus negatively regulate its biological activity. IL-18 is an IL-1-like cytokine that requires cleavage with caspase-1 to become active, was found to increase IgE production in a CD4+ T cells-, IL-4-, and STAT6-, dependent fashion. IL-18 and T cell receptor-mediated stimulation could induce naive CD4+ T cells to develop into IL-4-producing cells in vitro. Thus, caspase-1 and IL-18 may be critical in regulation of IgE production in vivo, providing a potential therapeutic target for allergic disorders. IL-18 production in primary synovial cultures and purified synovial fibroblasts was, in turn, upregulated by TNF- α , and IL-1 β , suggesting that monokine expression can feed back to promote Th1 cell development in synovial membrane. Besides, synergistic combinations of IL-18, IL-12, and IL-15 may be of importance in sustaining both Th1 responses and monokine production in RA.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine
Tel:400-999-2100

Email:techsupport@elabscience.cn

Web:www.elabscience.cn

Rev. V1.3