
## Recombinant Mouse SerpinB3/SCCA1 Protein (His Tag)

## Catalog Number: PKSM040578

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Mouse
Baculovirus-Insect Cells-derived Mouse SerpinB3/SCCA1 protein Met 1-Pro 386, with
an C-terminal His
46.5 kDa
46 & 95 kDa
NP_958751.2
Not validated for activity
>95 % as determined by reducing SDS-PAGE.
< 1.0 EU per µg of the protein as determined by the LAL method.
Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
This product is provided as lyophilized powder which is shipped with ice packs.
Lyophilized from sterile 50mM Tris 100mM NaCl, pH 8.0
Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
before lyophilization.
Please refer to the specific buffer information in the printed manual.
Please refer to the printed manual for detailed information.



> 95 % as determined by reducing SDS-PAGE.

## Background

Web:www.elabscience.com

## **Elabscience**®

Serpins are the largest and most diverse family of serine protease inhibitors which are involved in a number of fundamental biological processes such as blood coagulation, complement activation, fibrinolysis, angiogenesis, inflammation and tumor suppression and are expressed in a cell-specific manner. Serpins are a group of proteins with similar structures that were first identified as a set of proteins able to inhibit proteases. The acronym serpin was originally coined because many serpins inhibit chymotrypsin-like serine proteases (serine protease inhibitors). Over 1000 serpins have been identified. Mouse SerpinB3, also known as Squamous cell carcinoma antigen 1, SCCA-1, SERPINB3, SCCA and SCCA 1, is a cytoplasm protein which belongs to theserpin family and Ov-serpin subfamily. SerpinB3 may act as a protease inhibitor to modulate the host immune response against tumor cells. Mouse SerpinB3a and SerpinB3b, but not Serpinb3c, are functional, inhibiting both serine and cysteine proteinases with different inhibitory profiles due to the difference of two amino acids in their reactive site loops. SerpinB3a and SerpinB3b may play different roles by inhibiting intrinsic or extrinsic proteinases with different expression distributions and different inhibitory profiles.