Recombinant Human CXCL11 protein (His Tag)

Catalog Number: PDEH100884



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Mol_Mass	7.9 kDa
Accession	O14625
Bio-activity	Not validated for activity
Properties	
_	
Purity	> 95% as determined by reducing SDS-PAGE.
Purity Endotoxin	> 95% as determined by reducing SDS-PAGE. < 10 EU/mg of the protein as determined by the LAL method
·	, ,

 $^{\circ}$ C. Reconstituted protein solution can be stored at 4-8 $^{\circ}$ C for 2-7 days. Aliquots of reconstituted samples are stable at < -20 $^{\circ}$ C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CXCL11, also known as I-TAC, SCYB9B, H174 and beta-R1, is a non-ELR CXC chemokine. CXCL11 cDNA encodes a 94 amino acid (aa) residue precursor protein with a 21 aa residue putative signal sequence, which is cleaved to form the mature 73 aa residue protein. CXCL11 shares 36% and 37% amino acid sequence homology with IP-10 and MIG (two other known human non-ELR CXC chemokines), respectively. CXCL11 is expressed at low levels in normal tissues including thymus, spleen and pancreas. The expression of CXCL11 mRNA is radically up regulated in IFN-gamma and I L-1 stimulated astrocytes. Moderate increase in expression is also observed in stimulated monocytes. CXCL11 has potent chemoattractant activity for IL-2 activated T cells and transfected cell lines expressing CXCR3, but not freshly isolated T cells, neutrophils or monocytes. The gene encoding CXCL11 has been mapped to chromosome 4.

For Research Use Only