



A Reliable Research Partner in Life Science and Medicine

# **Mouse SAA Antibody Pair Set**

Catalog No. E-KAB-0716 Applications ELISA

**Synonyms** SAA1;PIG4;SAA;TP53I4;Serum amyloid A1

## Kit components & Storage

Title	Specifications	Storage
Mouse SAA Capture Antibody	1 vial, 100 µ g	Store at -20°C. Avoid freeze / thaw cycles.
Mouse SAA Detection Antibody (Biotin)	1 vial, 50 μL	Store at -20℃. Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0716)		
		Mayoo S A A Continue Antihody	Mouse SAA Detection Antibody	
		Mouse SAA Capture Antibody	(Biotin)	
Immunogen	Immunogen	Recombinant Mouse SAA protein	Recombinant Mouse SAA protein	
Information	Swissprot	P05366		
Product details	Reactivity	Mouse	Mouse	
	Host	Rat	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
			7.4	
	Purify	Protein A or G	Affinity purification	
	Specificity	Detects Mouse SAA in ELISAs.		

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>



A Reliable Research Partner in Life Science and Medicine

## **Applications**

Mouse SAA Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4ug/mL	Mouse SAA Capture	
Capture		Antibody	10 9
			Ajsu
ELISA	1:1000-1:10000	Mouse SAA Detection	Optical Density
	1.1000 1.10000		0.1
Detection		Antibody (Biotin)	
			0.01 1 10 100
			Mouse SAA concentration (ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

### **Background**

Serum amyloid A proteins (SAAs) are a family of homologous apolipoproteins of high density lipoprotein (HDL). They can be divided into two groups. The first group consists of the acute phase SAA1 and SAA2 that associate with HDL during inflammation and remodel the HDL particle by displacing apolipoprotein A1. The second group consists of constitutively expressed SAA4 and SAA5 that exist as minor apolipoproteins on HDL but make up more than 90% of the total SAA during homeostasis.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>