

# Recombinant Human PKM2 protein (His tag)

Catalog Number:PDEH100224



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	CTHBP;HEL-S-30;OIP3;PK3;PKM2;TCB;THBP1
<b>Species</b>	Human
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Ser 2-Pro 531
<b>Accession</b>	P14618
<b>Calculated Molecular Weight</b>	58.2 kDa
<b>Observed molecular weight</b>	59 kDa
<b>Tag</b>	N-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Pyruvate kinase isozymes M2 also known as pyruvate kinase muscle isozyme 2 (PKM2); pyruvate kinase type K; cytosolic thyroid hormone-binding protein (CTHBP); thyroid hormone-binding protein 1 (THBP1); or opa-interacting protein 3 (OIP3); is an isoenzyme of the glycolytic enzyme pyruvate kinase. Pyruvate kinase isozymes M2 / PKM2 is a protein involved in glycolysis. The encoded protein is a pyruvate kinase that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate to ADP; generating ATP and pyruvate. PKM2 has been shown to interact with thyroid hormone and may mediate cellular metabolic effects induced by thyroid hormones. PKM2 has been found to bind Opa protein; a bacterial outer membrane protein involved in gonococcal adherence to and invasion of human cells; suggesting a role of this protein in bacterial pathogenesis. Several alternatively spliced transcript variants encoding a few distinct isoforms

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human PKM2 protein (His tag)

Catalog Number:PDEH100224



have been reported. PKM2 functions as a glycolytic enzyme that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate (PEP) to ADP; generating ATP. PKM2 may stimulates POU5F1-mediated transcriptional activation. This protein Plays a general role in caspase independent cell death of tumor cells. The ratio between the highly active tetrameric form and nearly inactive dimeric form determines whether glucose carbons are channeled to biosynthetic processes or used for glycolytic ATP production. The transition between the 2 forms of PKM2 contributes to the control of glycolysis and is important for tumor cell proliferation and survival.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017