NFS1 Polyclonal Antibody

catalog number: E-AB-53070





Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|--|
| Reactivity | Human;Mouse |
| Immunogen | Fusion protein of human NFS1 |
| Host | Rabbit |
| Is otype | IgG |
| Purification | Antigen affinity purification |
| Conjugation | Unconjugated |
| buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

| Applications | Recommended Dilution |
|--------------|-----------------------------|
| WB | 1:1000-1:5000 |
| IHC | 1:100-1:200 |

Data





dilution of 1:100(×200)

Western blot analysis of HepG2 cell Hela cell Mouse kidney Immunohistochemistry of paraffin-embedded Human tissue K562 cell lysates using NFS1 Polyclonal Antibody at esophagus cancer tissue using NFS1 Polyclonal Antibody at dilution of 1:800 **Observed-MV:Refer to figures**

Calculated-MV:50 kDa

| Preparation & Storage | |
|-----------------------|---|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the |
| | temperature recommended. |

Background

Iron-sulfur clusters are required for the function of many cellular enzymes. The proteins encoded by this gene supply inorganic sulfur to these clusters by removing the sulfur from cysteine, creating alanine in the process. This gene uses alternate in-frame translation initiation sites to generate mitochondrial forms and cytoplasmic/nuclear forms. Selection of the alternative initiation sites is determined by the cytosolic pH. The encoded proteins belong to the class-V family of pyridoxal phosphate-dependent aminotransferases. Alternatively spliced transcript variants have been described.

For Research Use Only