
Recombinant Human DMP1 Protein (His Tag)

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Catalog Number: PKSH032347



Description	
•	н
Species	Human
Mol_Mass	55.0 kDa
Accession	Q13316
Bio-activity	Not validated for activity
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Histidine-HCl, 6% Trehalose, 4%
	Mannitol, 0.05% Tween 80, pH6.0.
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.



> 95 % as determined by reducing SDS-PAGE.

Background

Data

Dentin Matrix Acidic Phosphoprotein 1 (DMP-1) is an extracellular matrix protein and a member of the small integrin binding ligand N-linked glycoprotein family. DMP-1 is expressed in teeth particularly in odontoblast, ameloblast, and cementoblast. DMP-1 is critical for proper mineralization of bone and dentin. DMP-1 may have a dual function during osteoblast differentiation. In the nucleus of undifferentiated osteoblasts, the unphosphorylated form of DMP-1 acts as a transcriptional component for activation of osteoblast-specific genes like osteocalcin. During the osteoblast to osteocyte transition phase, DMP-1 is phosphorylated and exported into the extracellular matrix, where it regulates nucleation of hydroxyapatite. DMP-1 mutations have also been shown to cause rickets hypophosphatemic autosomal recessive type 1 (ARHR1).

For Research Use Only