

MEC/CCL28, Human, Recombinant

Cat. No. : PCK113

General Information

Synonyms	C-C Motif Chemokine 28;Mucosae-Associated Epithelial Chemokine;MEC;Protein CCK1; Small-Inducible Cytokine A28;CCL28;SCYA28
Species	Human
Expression host	E.coli
Sequence	Ser20-Tyr127
Accession	Q9NRJ3
Mol mass	12.49 kDa
Expiration date	12 months

Product feature

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin (EU/μg)	< 0.1
Storage	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
Shipping	Ice bag
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Background

Chemokine (C-C Motif) Ligand 28 (CCL28) is a novel Chemokine that shares the most homology with CCL27/CTACK. CCL28 shows chemotactic activity for resting CD4, CD8 T-cells and eosinophils. It Binds to CCR3 and CCR10 and induces calcium mobilization in a dose-dependent manner. CCR10 (GPR2 orphan Receptor) is also the Receptor for CCL27/CTACK. CCL28 is preferentially expressed by epithelial cells of diverse tissues, with highest expression level in normal and pathological colon. It is also expressed in normal and asthmatic lung tissues. Human and mouse CCL28 shares 83% sequence identity in their mature regions.

SDS-PAGE