Recombinant Mouse CD16-2/FCGR4 Protein (His Tag)

Catalog Number: PKSM041244

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Mouse
Source	HEK293 Cells-derived Mouse CD16-2/FCGR4 protein Gly21-Gln203, with an C-terminal
	His
Calculated MW	21.9 kDa
Observed MW	25-35 kDa
Accession	AAH27310.1
Bio-activity	Not validated for activity
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



Background

Elabscience®

Fcgr4, also known as CD16-2, is one of the receptors for Fc region of IgG which involve in immune responses. Fcgr4 mainly functions in cellular response to lipopolysaccharide, NK T cell proliferation, regulation of sensory perception of pain, wound healing etc. Three groups are included for Fc γ receptors (FcR), and they are Fc γ RI (CD64), Fc γ RII (CD3 2), and Fc γ RIII (CD16). Among these, CD64 possess high affinity even for monomeric IgG, while CD32 and CD16 display a relative lower affinity for IgG. Genes encodes these receptors are diverse differing by species and cell types. The aggregation of FcR having immunoreceptor tyrosine-based activation motifs (ITAMs) activates sequentially src family tyrosine kinases and syk family tyrosine kinases that connect transduced signals to common activation pathways shared with other receptors. FcR with ITAMs elicit cell activation, endocytosis, and phagocytosis. Fcgr4 belongs to Fc γ RIII (CD16) group.