Elabscience®

PKM2-specific Polyclonal Antibody

catalog number: E-AB-92125


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	A synthetic peptide of human PKM2-specific
Host	Rabbit
Is otype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000

1:500-1:2000 1:50-1:200


Data

IF




Western blot analysis of extracts of various cell lines using PKM2-specific Polyclonal Antibody at 1:1000 dilution.

> **Observed-MW:58 kDa** Calculated-MW:56 kDa/57 kDa/58 kDa



Immunofluorescence analysis of C6 cells using PKM2specific Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells using PKM2-Blue: DAPI for nuclear staining.

Immunofluorescence analysis of NIH-3T3 cells using specific Polyclonal Antibody at dilution of 1:100 (40x lens). PKM2-specific Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Elabscience®

This gene encodes a protein involved in glycolysis. The encoded protein is a pyruvate kinase that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate to ADP, generating ATP and pyruvate. This protein has been shown to interact with thyroid hormone and may mediate cellular metabolic effects induced by thyroid hormones. This protein has been found to bind Opa protein, a bacterial outer membrane protein involved in gonococcal adherence to and invasion of human cells, suggesting a role of this protein in bacterial pathogenesis. Several alternatively spliced transcript variants encoding a few distinct isoforms have been reported.

For Research Use Only