A Reliable Research Partner in Life Science and Medicine

Elab Fluor® Violet 450 Anti-Mouse CD25 Antibody[PC-61.5.3]

Catalog Number: E-AB-F1102Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

 Isotype
 Rat IgG1, κ

 Clone No.
 PC-61.5.3

Isotype Control Elab Fluor® Violet 450 Rat IgG1, κ Isotype Control[HRPN] [Product E-AB-F09822Q]

Conjugation Elab Fluor[®] Violet 450

Conjugation Information Elab Fluor[®] Violet 450 is designed to be excited by the violet laser (405 nm) and

detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein

protectant.


Applications

Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



C57BL/6 murine splenocytes are stained with Elab Fluor[®] Violet 450 Anti-Mouse CD25 Antibody and APC Anti-Mouse CD4 Antibody.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names IL-2 receptor subunit alpha;IL-2-RA;IL-2R subunit alpha;IL2-RA;IL2RA;Interleukin-2

receptor subunit alpha;TAC antigen;p55

 Uniprot ID
 P01590

 Gene ID
 16184

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Background

CD25 is a 55 kD glycoprotein, also known as the low affinity IL-2R α , Ly-43, p55, or Tac. It is expressed on activated T and B cells, thymocyte subset, pre-B cells, and T regulatory cells. In association with CD122 (IL-2R β) and CD132(common γ chain), CD25 forms the high affinity signaling IL-2 receptor.