

A Reliable Research Partner in Life Science and Medicine

APC Anti-Mouse CD161/NK1.1 Antibody[PK136]

Catalog Number: E-AB-F0987E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse
Host Mouse

Isotype Mouse IgG2a, κ

Clone No. PK136

Isotype Control APC Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802E]

Conjugation APC

Conjugation Information APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine splenocytes with FITC Anti-Mouse CD49b/pan-NK cells Antibody[DX5] and APC Anti-Mouse CD161/NK1.1 Antibody[PK136] (left) or APC Mouse IgG2a, κ Isotype Control (right). Total viable cells were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping lce bag

Antigen Information

Alternate Names CD161 antigen-like family member C;CD161;NK1.1;CD161c;Killer cell lectin-like

receptor subfamily B member 1C;Klrb1c;Ly-55c;NKR-P1 40;NKR-P1.9;NKR-P1C

Uniprot ID P27814;P27812;Q99JB4

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Gene ID Background 17059;80782

NK-1.1 surface antigen, also known as CD161b/CD161c and Ly-55, is encoded by the NKR-P1B/NKR-P1C gene. It is expressed on NK cells and NK-T cells in some mouse strains, including C57BL/6, FVB/N, and NZB, but not AKR, BALB/c, CBA/J, C3H, DBA/1, DBA/2, NOD, SJL, and 129. Expression of NKR-P1C antigen has been correlated with lysis of tumor cells in vitro and rejection of bone marrow allografts in vivo. NK-1.1 has also been shown to play a role in NK cell activation, IFN-γ production, and cytotoxic granule release. NK-1.1 and DX5 are commonly used as mouse NK cell markers.