A Reliable Research Partner in Life Science and Medicine

Catenin beta Monoclonal Antibody

catalog number: E-AB-22111

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat **Immunogen** Recombinant Protein

Host Mouse Isotype IgG Clone 2E4

Purification Protein A purification


Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein Buffer

protectant and 50% glycerol.


Applications Recommended Dilution

WB 1:500-1:2000 1:100-1:300 IHC IF 1:100-1:300

Data




Rat liver with Catenin beta Monoclonal Antibody



Western Blot analysis of 1) Hela, 2) 293T, 3) Mouse liver, 4) Immunohistochemistry of paraffin-embedded Mouse kidney tissue using Catenin beta Monoclonal Antibody at dilution of 1:200.





Immunofluorescence analysis of Human stomach cancer tissue using Catenin beta Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Storage

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Key dowstream component of the canonical Wnt signaling pathway. In the absence of Wnt, forms a complex with AXIN1, AXIN2, APC, CSNK1A1 and GSK3B that promotes phosphorylation on N-terminal Ser and Thr residues and ubiquitination of CTNNB1 via BTRC and its subsequent degradation by the proteasome. In the presence of Wnt ligand, CTNNB1 is not ubiquitinated and accumulates in the nucleus, where it acts as a coactivator for transcription factors of the TCF/LEF family, leading to activate Wnt responsive genes. Involved in the regulation of cell adhesion. The majority of beta-catenin is localized to the cell membrane and is part of E-cadherin/catenin adhesion complexes which are proposed to couple cadherins to the actin cytoskeleton.

For Research Use Only

Toll-free: 1-888-852-8623 Web:w w w .elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com