

IL-16/LCF, Mouse, Recombinant

Cat. No. : PCK322

General Information

Synonyms	LCF (Lymphocyte Chemoattractant Factor);mKIAA4048
Species	Mouse
Expression host	E.coli
Sequence	MHDLNSSTD SAASASAASDISVESKEATVCTVTLEKTSAGLGFSLEGGKGS LHGDKPLTINRIF KGDRTGEMVQPGDEILQLAGTAVQGLTRFEAWNVIKALPDGPVTIVIRRTSLQCKQTTASADS with polyhistidine tag at the C-terminus.
Accession	AAC04383.1
Tag	His-tag at the C-terminus
Mol mass	14.04 kDa
Expiration date	12 months

Product feature

Purity	> 95% as determined by SDS-PAGE. Ni-NTA chromatography.
Endotoxin (EU/μg)	< 0.1
Storage	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
Shipping	Ice bag
Formulation	The protein was lyophilized from a 0.2 μm filtered solution containing 1 × PBS, pH 7.4.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Background

Interleukin 16 (IL-16) is a cytokine that is released by a variety of cells (including lymphocytes and some epithelial cells) that has been characterized as a chemoattractant for certain immune cells expressing the cell surface molecule CD4. The protein encoded by this gene is a pleiotropic cytokine that functions as a chemoattractant, a modulator of T cell activation, and an inhibitor of HIV replication. The signaling process of this cytokine is mediated by CD4. The product of this gene undergoes proteolytic processing, which is found to yield two functional proteins.

SDS-PAGE

