



MAP2K6 Polyclonal Antibody

catalog number: E-AB-40167

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|--|
| Reactivity | Mouse;Rat |
| Immunogen | Recombinant Mouse Dual specificity mitogen-activated protein kinase kinase 6 protein |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Antigen Affinity Purification |
| Conjugation | Unconjugated |
| Buffer | PBS with 0.05% Proclin300, 1% protective protein and 50% glycerol, pH7.4 |

| Applications | Recommended Dilution |
|--------------|----------------------|
| IHC | 1:50-1:100 |

| Data | |
|--|--|
|  |  |
| Immunohistochemistry of paraffin-embedded Mouse brain using MAP2K6 Polyclonal Antibody at dilution of 1:50 | Immunohistochemistry of paraffin-embedded Rat brain using MAP2K6 Polyclonal Antibody at dilution of 1:50 |

| Preparation & Storage | |
|-----------------------|--|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

| Background | |
|---|--|
| <p>This gene encodes a member of the dual specificity protein kinase family, which functions as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein phosphorylates and activates p38 MAP kinase in response to inflammatory cytokines or environmental stress. As an essential component of p38 MAP kinase mediated signal transduction pathway, this gene is involved in many cellular processes such as stress induced cell cycle arrest, transcription activation and apoptosis.</p> | |