

## Recombinant Mouse NOV/CCN3 Protein (His Tag)

**Catalog Number: PKSM041114**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	HEK293 Cells-derived Mouse NOV/CCN3 protein Ser26-Ile354, with an C-terminal His
<b>Calculated MW</b>	37.1 kDa
<b>Observed MW</b>	50 kDa
<b>Accession</b>	Q64299
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

NOV, also called CCN3, is a secreted protein of CCN family members. CCN family members are highly conserved cysteine rich proteins sharing a common modular structure having 4 conserved domains, insulin-like growth factor-binding protein (IGFBP) domain, von Willebrand type C (VWC) domain, thrombospondin-1 (TSP-1) domain, and C-terminal (CT) domain (absent in CCN5). By specific interactions with these domains, CCN proteins modulate multiple signalling pathways including BMPs, Wnt, TGFs, Notch and integrins to regulate cell proliferation, differentiation, adhesion, migration, angiogenesis, and survival. CCN3 is firstly characterized as a promoter of progenitor activity of human hematopoietic stem cells, as knockdown of CCN3 can abrogate the function of primitive progenitors. Recent studies showed that CCN3 is also actively involved in the process of wound healing. CCN3 is highly expressed in granulation tissues of cutaneous wounds and capable of inducing synthetic responses of fibroblasts.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017