

## Mitochondrial Extraction Assay Kit

**Catalog No:** E-BC-E001

**Specification:** 50 Assays/100 Assays

- Note:**
- ① All steps of mitochondria isolation were performed on ice box or at 4 °C, and the solutions should be placed in ice water for pre-cooling before use.
  - ② Use a glass homogenizer with tight clearance.

### General information

**Intended use** This kit can be used to extract mitochondrial from animal tissue.

**Detection principle** The principle of mitochondrial separation and purification: after rupturing cells mechanically, the debris and huge organelles are removed by low-speed differential centrifugation, then purified mitochondria will be obtained by high-speed differential centrifugation.

### Kit components & storage

Item	Component	Size 1 (50 Assays)	Size 1 (100 Assays)	Storage
Reagent 1	Lysis Buffer	30 mL × 1 vial	60 mL × 1 vial	-20 °C, 12 months
Reagent 2	Mito-Wash Buffer	15 mL × 1 vial	30 mL × 1 vial	-20 °C, 12 months, shading light
Reagent 3	Store Buffer	7 mL × 1 vial	15 mL × 1 vial	-20 °C, 12 months

**Note:** The reagents must be stored strictly according to the preservation conditions in the above table.

The reagents in different kits cannot be mixed with each other.

### Materials prepared by users

**Instruments:**

High-speed freezing centrifuge, 5 mL Glass homogenizer

**Reagents:**

Double distilled water, PBS(0.01 M, pH 7.4)

### Reagent preparation

Place all reagents in ice water for pre-cooling for at least 15 min after they have returned to the solution state before use.

#### For Research Use Only

Thank you for your recent purchase.  
If you would like to learn more about biochemical kit, please visit [www.elabscience.com](http://www.elabscience.com)

Focus on lab research  
Service for life science

## Operation table

- ① Weight 0.05 g~0.1 g of fresh tissue and wash with PBS (0.01 M, pH 7.4) at 2-8℃ to remove blood. Blot the water with absorbent paper. Cut the tissue into pieces with scissors and place them into a pre-cooled 5 mL glass homogenizer. Add 0.5 mL of pre-cooled lysis buffer. Grind the tissue up and down in the ice bath for about 30-40 times.
- ② Transfere tissue homogenate to a 2 mL pre-cooled EP tube, centrifuge at 500×g at 4 ℃ for 5 min.
- ③ Transferre the supernatant to a new 2 mL pre-cooled EP tube, centrifuge at 500×g at 4 ℃ for 5 min.
- ④ Transferre the supernatant to a new 2 mL pre-cooled EP tube, centrifuge at 15000×g at 4 ℃ for 15 min.
- ⑤ Discard the supernatant, resuspend the precipitate with 0.2 mL of pre-cooled mito-wash buffer, then centrifuge with 11000×g at 4 ℃ for 10 min.
- ⑥ Discard the supernatant, resuspend the precipitate with 50-100 μL of pre-cooled store buffer. The prepared mitochondrial extract can be used immediately or be stored at -70 ℃, avoiding repeated freeze-thaw.