## Recombinant Mouse IFN alpha 1a protein(His Tag)

Catalog Number: PKSM041492



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Mouse
Mol_Mass	19.9 kDa
Accession	P01572
Bio-activity	Measure by its ability to protect L929 cells infected with encephalomyocarditis (EMC)
	virus. The ED <sub>50</sub> for this effect is <10 pg/mL. The specific activity of recombinant
	mouse IFN alpha 1a is $> 4 \times 10^7$ IU/mg.
Properties	
Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.1 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

This product is provided as lyophilized powder which is shipped with ice packs.

reconstituted samples are stable at < -20°C for 3 months.

**Formulation** Lyophilized from sterile PBS, pH 7.4.

Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants


before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data

Shipping



> 98 % as determined by reducing SDS-PAGE.

## Background

IFNA1, also known as IFN-alpha and IFNA, belongs to the alpha/beta interferon family. Interferons(IFNs) are proteins made and released by host cells in response to the presence of pathogens such as viruses, bacteria, parasites, or tumor cells. They belong to the large class of glycoproteins known as cytokines. IFNs stimulate the production of two enzymes: a protein kinase and an oligoadenylate synthetase. They allow for communication between cells to trigger the protective defenses of the immune system that eradicate pathogens or tumors. IFNs can activate immune cells, such as natural killer cells and macrophages; they increase recognition of infection or tumor cells by up-regulating antigen presentation to T lymphocytes, and they also increase the ability of uninfected host cells to resist new infection by the virus. Leukocyte interferon is produced predominantly by B lymphocytes. Immune interferon is produced by mitogen-or antigen-stimulated T lymphocytes. IFNA1 is produced by macrophages and has antiviral activities.

## For Research Use Only