# **Elabscience**®

## **EIF5B** Polyclonal Antibody

### catalog number: E-AB-66431

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description  |  |
|--------------|--|
| Reactivity   | Human;Mouse;Rat  |
| Immunogen    | Recombinant fusion protein of human EIF5B (NP_056988.3).                           |
| Host         | Rabbit   |
| Isotype      | IgG  |
| Purification | Affinity purification  |
| Buffer       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
|              | De commende d'Dilation   |

| Applications | Recommended Dilution |
|--------------|----------------------|
| WB           | 1:500-1:2000         |
| IF           | 1:50-1:200           |

#### Data





Western blot analysis of extracts of various cell lines using

### EIF5B Polyclonal Antibody. Observed-MW:179 kDa

#### Calculated-MW:138 kDa



Immunofluorescence analysis of C6 cells using EIF5B Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using EIF5B Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining. Immunofluorescence analysis of U-2 OS cells using EIF5B Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

| Preparation & Storage |  |
|-----------------------|--|
| Storage               | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.  |
| Shipping              | The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended. |

#### Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com

# **Elabscience**®

Accurate initiation of translation in eukaryotes is complex and requires many factors, some of which are composed of multiple subunits. The process is simpler in prokaryotes which have only three initiation factors (IF1, IF2, IF3). Two of these factors are conserved in eukaryotes: the homolog of IF1 is eIF1A and the homolog of IF2 is eIF5B. This gene encodes eIF5B. Factors eIF1A and eIF5B interact on the ribosome along with other initiation factors and GTP to position the initiation methionine tRNA on the start codon of the mRNA so that translation initiates accurately.

For Research Use Only