

Recombinant Mouse IL6ST/CD130 Protein (His Tag)

Catalog Number: PKSM040841

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse IL6ST/CD130 protein Met 1-Glu 617, with an C-terminal His
Calculated MW	68.0 kDa
Observed MW	80-90 kDa
Accession	NP_034690.3
Bio-activity	Measured by its ability to inhibit the IL-6R α enhancement of IL-6 activity on M1 mouse myeloid leukemia cells. The ED ₅₀ for this effect is typically 0.6-3 μ g/mL in the presence of 50 ng/mL recombinant human IL-6sR and 100 ng/mL recombinant human IL-6.


Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.



Measured by its ability to inhibit the IL-6R α enhancement of IL-6 activity on M1 mouse myeloid leukemia cells. The ED₅₀ for this effect is typically 0.6-3 μ g/mL in the presence of 50 ng/mL recombinant human IL-6sR and 100 ng/mL recombinant human IL-6.

Background

For Research Use Only

Glycoprotein 130 (also known as gp130, IL6ST, IL6-beta or CD130) is a transmembrane protein which is the founding member of the class of all cytokine receptors. CD130/gp130 is a signal transducer shared by many cytokines, including interleukin 6 (IL6), ciliary neurotrophic factor (CNTF), leukemia inhibitory factor (LIF), and Oncostatin M (OSM). CD130/gp130 functions as a part of the cytokine receptor complex. The activation of this protein is dependent upon the binding of cytokines to their receptors. CD130/gp130 plays a critical role in regulating myocyte apoptosis. Alternatively spliced transcript variants encoding distinct isoforms have been described. A related pseudogene has been identified on chromosome 17. The receptor systems for IL6, LIF, OSM, CNTF, IL11, CTF1 and BSF3 can utilize gp130 for initiating signal transmission. CD130/gp130 binds to IL6/IL6R (alpha chain) complex, resulting in the formation of high-affinity IL6 binding sites, and transduces the signal. CD130/gp130 may have a role in embryonic development. The type I OSM receptor is capable of transducing OSM-specific signaling events.