

Recombinant Mouse PTK6/Brk Protein (His & GST Tag)

Catalog Number: PKSM040298

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	Baculovirus-Insect Cells-derived Mouse PTK6/Brk protein Met1-Val451, with an N-terminal His & GST
Calculated MW	79.8 kDa
Observed MW	66 kDa
Accession	Q64434
Bio-activity	The specific activity was determined to be 5 nmol/min/mg using poly [Glu, Tyr] 4:1 as substrate.

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Concentration	Subject to label value.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20°C.
Formulation	Supplied as sterile solution of 20mM Tris, 500mM NaCl, pH 7.4, 10% glycerol

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Tyrosine kinase (PTKs) is a protein that carry out tyrosine phosphorylation, which play a fundamental role in cell proliferation, survival, adhesion, and motility and have also been demenstrated to mediate malignant cell transformation. Overexpression of this protein in mammary epithelial cells leads to sensitization of the cells to epidermal growth factor and results in a partially transformed phenotype. Two classes of PTKs are present in cells: the transmembrane receptor PTKs and the non-receptor PTKs. Tyrosine kinase(PTKs)-6/ BRK is a cytoplasmic non-receptor protein kinase which may function as an intracellular signal transducer in epithelial tissues. Tyrosine kinase(PTKs)-6/ BRK has been shown to undergo autophosphorylation. It has been found that the constitutive expression of the tyrosine kinase(PTKs)-6/ BRK is in a large proportion of cutaneous T-cell lymphomas and other transformed T- and B-cell populations. State BRK expression was also induced in normal T-cells. In clinical, the cytoplasmic tyrosine kinase PTK6 (BRK) shows elevated expression in approximately two-thirds of primary breast tumours, and is implicated in EGF receptor-dependent signalling and epithelial tumorigenesis.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017