

A Reliable Research Partner in Life Science and Medicine

RAD50 Polyclonal Antibody

catalog number: E-AB-63998

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse

Immunogen A synthetic peptide of human RAD50

Host Rabbit Isotype IgG


Purification Affinity purification


Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications Recommended Dilution

WB 1:500-1:2000 **IHC** 1:50-1:100

Data







Western blot analysis of extracts of various cell lines using RAD50 Polyclonal Antibody at dilution of 1:500.

Observed-MV:180 kDa

Calculated-MV: 138 kDa/153 kDa/154 kDa



Immunohistochemistry of paraffin-embedded Human breast cancer using RAD50 Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded Human stomach using RAD50 Polyclonal Antibody at dilution of 1:100 (40x lens).

Immunohistochemistry of paraffin-embedded Human gastric cancer using RAD50 Polyclonal Antibody at dilution of 1:100 (40x lens).

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack,upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

The protein encoded by this gene is highly similar to Saccharomyces cerevisiae Rad50, a protein involved in DNA double-strand break repair. This protein forms a complex with MRE11 and NBS1. The protein complex binds to DNA and displays numerous enzymatic activities that are required for nonhomologous joining of DNA ends. This protein, cooperating with its partners, is important for DNA double-strand break repair, cell cycle checkpoint activation, telomere maintenance, and meiotic recombination. Knockout studies of the mouse homolog suggest this gene is essential for cell growth and viability. Mutations in this gene are the cause of Nijmegen breakage syndrome-like disorder.

Fax: 1-832-243-6017