
Recombinant Human CLDN1 protein(His Tag)

Catalog Number: PDEH100757

Note: Centrifuge before opening to ensure complete recovery of vial contents.

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Description	
Calculated MW9.6 kDaObserved MW14 kDaAccession095832Bio-activityNot validated for activityPropertiesPurity> 90% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Species	Human
Observed MW14 kDaAccessionO95832Bio-activityNot validated for activityPropertiesPurity> 90% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Source	E.coli-derived Human CLDN1 protein Gln29-Lys115, with an N-terminal His
AccessionO95832Bio-activityNot validated for activityPropertiesPurity> 90% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Calculated MW	9.6 kDa
Bio-activityNot validated for activityPropertiesPurity> 90% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Observed MW	14 kDa
PropertiesPurity> 90% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL methodStorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to - °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.	Accession	O95832
Purity> 90% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Bio-activity	Not validated for activity
Endotoxin< 10 EU/mg of the protein as determined by the LAL methodStorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to - °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.	Properties	
StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to - °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Purity	> 90% as determined by reducing SDS-PAGE.
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Shippingreconstituted samples are stable at < -20°C for 3 months.	Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.		°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.		reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Mannitol.	Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
	Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with 5% Trehalose and 5%
Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution		Mannitol.
it is reconstruction and store state of the value of the	Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of
0.5 mg/mL. Concentration is measured by UV-Vis.		0.5 mg/mL. Concentration is measured by UV-Vis.





> 90 % as determined by reducing SDS-PAGE.

Background

Claudins function as major constituents of the tight junction complexes that regulate the permeability of epithelia. While some claudin family members play essential roles in the formation of impermeable barriers, others mediate the permeability to ions and small molecules. Often, several claudin family members are coexpressed and interact with each other, and this determines the overall permeability. CLDN1 is required to prevent the paracellular diffusion of small molecules through tight junctions in the epidermis and is required for the normal barrier function of the skin. Required for normal water homeostasis and to prevent excessive water loss through the skin, probably via an indirect effect on the expression levels of other proteins, since CLDN1 itself seems to be dispensable for water barrier formation in keratinocyte tight junctions.

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>