## Recombinant Human PDCD1/PD-1/CD279 (C-6His-Avi) Biotinylated

Catalog Number: PKSH033853



Description		
Species	Human	
Mol_Mass	19.2 kDa	
Accession	Q15116	
Bio-activity	Not validated for activity	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of	
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.	
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants	
	before lyophilization.	
	Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	

Note: Centrifuge before opening to ensure complete recovery of vial contents.

kDa	МК	R
120 90		and the second
60		S. Starley
40		
30	-	
20		-
14		
14	eccediti	

> 95 % as determined by reducing SDS-PAGE.

## Background

Programmed cell death protein 1(PDCD1) is a single-pass type I membrane protein and contains 1 Ig-like V-type domain. PD-1 is a member of the extended CD28/CTLA-4 family of T cell regulators. PDCD1 inhibits the T-cell proliferation and production of related cytokines including IL-1, IL-4, IL-10 and IFN-γ by suppressing the activation and transduction of PI3K/AKT pathway. In addition, coligation of PDCD1 inhibits BCR-mediating signal by dephosphorylating key signal transducer. PDCD1 has been suggested to be involved in lymphocyte clonal selection and peripheral tolerance, and thus contributes to the prevention of autoimmune diseases. As a cell surface molecule, PDCD1 regulates the adaptive immune response. Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function.

## For Research Use Only